



STOVEK
veterinary production




Antibiotic powders

AMOCCLAV amoxicillin 50%, clavulanic acid 12,5%	●				4
AMOCOL 50 amoxicillin 50%, colistin 4 000 000 IU/gram	●	●			4
AMOXICILLIN 80-TRV amoxicillin 80%	●	●			4
COLI 12-TRV colistin 12 000 000 IU/gram	●	●			4
DOXYGIL 50 doxycycline 50%	●	●			5
DOXYTIL doxycycline 10%, tylosin 10%	●	●			5
ENROFLOXACIN 20-TRV enrofloxacin 20%	●				5
NOSIGEPTID 1%-TRV nosigeptide 1%	●	●			5
OXYTETRACYCLINE 100-TRV oxytetracycline 90%	●	●	●		6
TILTAR 80 tylosin 80%	●				6

Coccidiostatics

AMPROLIUM 100-TRV amprolium 90%	●	●		●	6
SALINOMYCIN 12-TRV salinomycin 12%	●	●		●	6

Antibiotic solutions

CIPROFLOXACIN AQUA ciprofloxacin 20%	●				7
COLISTIN AQUA colistin 4 000 000 IU/ml	●	●			7
DOXICOLIN AQUA colistin 1 000 000 IU/ml doxycycline 20%	●				7
ENROFLOXACIN 20% AQUA-TRV enrofloxacin 20%	●	●			7
ENROGENT AQUA-TRV enrofloxacin 10%, gentamicin 5%	●				8
ENROMOL AQUA-TRV enrofloxacin 10%, colistin 2 000 000 IU/ml	●	●			8
ENROTRIM AQUA enrofloxacin 10%, trimethoprim 5%	●				8
FLOROL 10% AQUA-TRV florfenicol 10%	●				8
LESTIN AQUA-TRV levofloxacin 10%, colistin 1 000 000 IU/ml	●				9
LEXOL AQUA levofloxacin 15%	●				9
SULFAPRIM AQUA-TRV sulfadimidine 20%, trimethoprim 4%	●	●			9
 TETRAFLOX AQUA-TRV tilmicosine 5%, levofloxacin 2,5% trimetoprim 2,5%	●				9
TILMICOSIN 30% AQUA-TRV tilmicosine 30%	●	●			10
TRIENROKOL AQUA enrofloxacin 10%, trimethoprim 5%, colistin 1 000 000 IU/ml	●	●			10
TYLOSIN AQUA tylosin 20%	●				10

Injection antibiotics

AMOXIDGEKT LA amoxicillin 15%		●			11
AZITHROMYCIN 10%-TRV azithromycin 10%		●		●	11
CEFTIDGECT LA-TRV ceftiofur 10%	●	●		●	11
ENRODGECT 100 enrofloxacin 10%		●		●	11



Injection antibiotics

FLOREZOL florfenicol 40%		●		●	12
GENTALUX gentamicin 4%		●			12
LEXODGEKT levofloxacin 15%		●		●	12
MOXIDGEKT LA-TRV moxifloxacin 10%		●			12
OXICET-TRV oxytetracycline 20%, ketoprofen 3%		●		●	13
OXYTETRACYCLINE 200 LA-TRV oxytetracycline 20%		●		●	13
NEW PENSTREP LA-TRV benzylpenicillin procaine 10% benzylpenicillin benzathine 10% dihydrostreptomycin sulfate 20%		●		●	13
TULATHROMYCIN 10-TRV tulatromycin 10%		●			13
TYLDGEKT tylosin 20%		●		●	14

Hormonal drugs

DECSA-TRV dexamethasone 0,4%		●	●	●	14
CLOPROSTENOL-TRV cloprostenol 0,025%		●			14
OXYTOCIN 10 ME-TRV oxytocin 10 IU/ml		●		●	14

Anti-inflammatory drugs

KETOPROFEN 10-TRV ketoprofen 10%		●	●		15
------------------------------------	--	---	---	--	----

Metabolic stimulators

FERRUM 20/B12-TRV iron 20%, vitamin B12 0,02%		●			15
---	--	---	--	--	----

Antiparasitic drugs

IVERMECTIN 1%-TRV ivermectin 1%		●		●	15
NEW IVERMECTIN AQUA ivermectin 4%	●	●		●	15

Antiviral drugs

TRIVIRON FORTE-TRV aRNase 3%	●	●			16
TRIVIDGEKT 5%-TRV aRNase 5%		●			16

Feed additives

NEW LYSINOFID-TRV L-lysine 95%	●	●		●	16
NEW TREFID-TRV L-threonine 95%	●	●		●	16

Probiotics

PROVAGEN B. Subtilis, B. licheniformis 10 ⁹ CFU/gram	●	●	●		17
PROVAGEN CONCENTRATE B. Subtilis, B. licheniformis 10 ¹¹ CFU/gram	●	●	●		17
NEW TRIOBAKT B. licheniformis 5x10 ⁹ CFU/gram	●				17

Biodestructor of manure

BIOSANVIT-M Bacillus 10 ⁷ CFU/gram	●	●			17
---	---	---	--	--	----

Antibiotic powders


AMOCLAV

amoxicillin 50%
clavulanic acid 12,5%



Indications for use:

Amoklav is used as a therapeutic agent for poultry infections of the gastrointestinal, respiratory tract and urinary tract caused by microorganisms sensitive to amoxicillin.

Dosage: {  Poultry - 50-100 gram/ton water

Storage conditions:

Amoclav is stored according to the requirement of the list B in a dry, dark place at a temperature from 0 °C to 25 °C. The shelf life is three years from the date of manufacture, subject to the storage terms.



AMOCOL 50

amoxicillin 50%
colistin 4 000 000 IU/gram



Indications for use:

Amocol 50 is used as a therapeutic agent for calves, and poultry with diseases, the causative agents of which are sensitive to the components of the veterinary drug.

Dosage: {  Poultry - 300 gram/ton water
 Calves - 3 gram/ 100 kg of body weight

Storage conditions:

Amocol 50 is stored according to the requirements of the list B in a dry, dark place at a temperature from 0 °C to 25 °C. The shelf life is three years from the date of manufacture, subject to the storage terms.



AMOXICILLIN 80-TRV

amoxicillin 80%



Indications for use:

Amoxicillin 80-TRV is used in calves and poultry with a therapeutic purpose for pneumoniae disease bacterial etiology, the causative agents of which are sensitive to amoxicillin.

Dosage: {  Poultry - 100-250 gram/ton water
 Calves - 15-30 mg/ 1 kg of body weight

Storage conditions:

Amoxicillinum 80-TRV is stored according to list B in a dry, dark place at a temperature from 0 °C to 25 °C. The shelf life is three years from the date of manufacture, subject to the storage terms.


COLI 12-TRV

colistin 12 000 000 IU/g



Indications for use:

Coli 12-TPV is used to calves and poultry with colibacteriosis, salmonellosis and other infectious diseases of the gastrointestinal tract of bacterial etiology, the pathogens of which are sensitive to colistin.

Dosage: {  Poultry - 50-180 gram/ton water
 Calves - 5-8 mg/kg of body weight

Storage conditions:

Coli 12-TRV is stored under the requirement of the list B: in a dry, dark place at temperatures from 0°C to 25°C. Shelf life is three years from the date of manufacture, subject to storage conditions.

Antibiotic powders

DOXYGIL 50



doxycycline 50%



Indications for use:

Doxygil 50 is used as a therapeutic agent in calves and poultry for colibacillosis, salmonellosis, pasteurellosis, mycoplasmosis and other infectious diseases of the gastrointestinal tract, respiratory system, caused by microorganisms sensitive to doxycycline.

Dosage:

-  Poultry - 20-40 mg/1 kg of body weight
-  Calves - 0,1 gram/ 10 kg of body weight

Storage conditions:

Doxygil 50 is stored according to list B in a dry, dark place at a temperature from 0 °C to 25 °C. Shelf life - 3 (three) years from the date of manufacture, subject to storage and transportation conditions.

DOXYTIL



doxycycline 10%
tylosin 10%



Indications for use:

Doxytil is a semisynthetic tetracycline derivative. Active against gram-positive and gram-negative microorganisms, incl. Actinobacillus pleuropneumoniae, Clostridium spp., Escherichia coli, Staphylococcus spp., Streptococcus spp., Haemophilus spp., Pasteurella spp.

Dosage:

-  Poultry - 0,5-1 kg/ton water
-  Calves - 5 gram/ 100 kg of body weight

Storage conditions:

Doxytil is stored according to list B in a dry, dark place at a temperature from 0 °C to 25 °C. Shelf life - three years from the date of manufacture, subject to storage conditions.

ENROFLOXACIN 20-TRV


enrofloxacin 20%



Indications for use:

Enrofloxacin 20-TPV is used as a therapeutic agent in poultry for respiratory and gastrointestinal diseases of bacterial etiology: colibacillosis, salmonellosis, pasteurellosis, mycoplasmosis, as well as coinfections and secondary infections, the causative agents of which are sensitive to enrofloxacin.

Dosage:

-  Poultry - 500 gram/ton of water or feed

Storage conditions:

Enrofloxacin 20-TPV is stored according to list B in a dry, dark place at a temperature from 0 °C to 25 °C. Shelf life - three years from the date of manufacture, subject to storage and transportation conditions.

NOSIHEPTIDE 1%-TRV



nosiheptide 1%



Indications for use:

Nosiheptide 1%-TRV is used for therapeutic purposes in cattle and poultry with clostridium, echirichiosis, salmonellosis, streptococcosis and other bacterial diseases caused by microorganisms, sensitive to nozigeptide.

Dosage:

-  Poultry - 200-250 gram/ton feed
-  Calves - 300-500 gram/ ton feed

Storage conditions:

Nosiheptide 1%-TRV is stored according to list B in a dry, dark place at a temperature from 0 °C to 25 °C. Shelf life - two years from the date of manufacture, subject to storage and transportation conditions.

Antibiotic powders

OXYTETRACYCLIN 100-TRV





oxytetracycline 90%

Indications for use:

Oxytetracyclin 100-TRV is used in cattle, poultry, horses as a therapeutic agent for diseases of the respiratory, genitourinary system, gastrointestinal tract, secondary diseases of bacterial etiology.

Dosage:

-  Poultry - 20-40 mg/kg of body weight
-  Cattle - 10-20 mg/kg of body weight

Storage conditions:

Oxytetracyclin 100-TRV is stored according to list B in a dry, dark place at a temperature from 0 °C to 25 °C. The shelf life in the manufacturer's packaging, subject to the shelf life of three years from the date of manufacture.

TILTAR 80




tylosin 80%

Indications for use:

Tiltar 80 is used as a therapeutic agent for respiratory mycoplasmosis of chickens, infectious sinusitis of turkeys.

Dosage:

-  Poultry - 0,5 kg/ton water

Storage conditions:

Tiltar 80 is stored under the requirement of the list B: in a dark place at temperatures from 0°C to 25°C. Shelf life is three years from the date of manufacture, subject to storage conditions.

Coccidiostatics

AMPROLIUM 100-TRV






amprolium 90%

Indications for use:

Amprolium 100-TRV is used as a therapeutic and prophylactic agent for eimeriosis in broiler chickens, replacement poultry, breeding poultry, as well as cattle and small cattle.

Dosage:

-  Poultry - 130-300 gram/ton of water or feed
-  Calves - 130-300 gram/ton of water or feed
-  Sheep - 130-300 gram/ton of feed

Storage conditions:

Amprolium 100-TRV is stored in a dry, dark place at a temperature from 0 °C to 25 °C. Shelf life - three years from the date of manufacture, subject to storage conditions.

SALINOMYCIN 12-TRV





salinomycin 12%

Indications for use:

Salinomycin 12-TRV is an anti-eimeriosis agent of a broad spectrum of action of the ionophore group. It has a coccidiostatic effect in the early stages of development of coccidia.

Dosage:

-  Poultry - 500 gram/ton of feed
-  Calves - 100-250 gram/ton of feed

Storage conditions:

Salinomycin 12-TRV is stored in a dry, dark place at a temperature from 0 °C to 25 °C. The shelf life in the manufacturer's packaging, subject to the shelf life of three years from the date of manufacture.


CIPROFLOXACIN AQUA

ciprofloxacin 20%



Indications for use:

Ciprofloxacin aqua is used as a therapeutic agent in poultry for treatment of the gastrointestinal and respiratory tract, the genitourinary system and other diseases caused by microorganisms, sensitive to ciprofloxacin.

Dosage: {  Poultry - 250-330 ml/ton water

Storage conditions:

Ciprofloxacin aqua is stored with precautions according to list B in a dry, dark place at a temperature from 5 °C to 25 °C. The shelf life, if the storage conditions are observed in the sealed manufacturer's packaging, is two years from the date of manufacture.



COLISTIN AQUA

colistin 4 000 000 IU/ml



Indications for use:

The veterinary product is used to treat calves and poultry with bacterial diseases of the gastrointestinal tract, the causative agents of which are sensitive to colistin.

Dosage: {  Poultry - 100-125 ml/ton water
 Calves - 0,01-0,025 ml/kg of body weight

Storage conditions:

Colistin aqua is stored under the requirement of the list B: in a dark place at temperatures from 5°C to 25°C. Shelf life is two years from the date of manufacture.


DOXICOLIN AQUA

doxycycline 20%
colistin 1 000 000 IU/ml



Indications for use:

Doxicolin aqua is used as a therapeutic agent in poultry for diseases of salmonellosis caused by microorganisms sensitive to colistin and doxycycline.

Dosage: {  Poultry - 0,5 l/ton water

Storage conditions:

Doxicolin aqua is stored according to list B in a dry, dark place at a temperature from 5 °C to 25 °C. Shelf life is two years from the date of manufacture, subject to storage conditions.



ENROFLOXACIN 20% AQUA-TRV

enrofloxacin 20%



Indications for use:

Enrofloxacin 20% aqua-TRV is used as a therapeutic agent in cattle, poultry for respiratory and gastrointestinal diseases of bacterial etiology.

Dosage: {  Poultry - 0,25-0,5 l/ton water
 Calves - 1,25-2,5 ml/100 kg of body weight

Storage conditions:

The veterinary drug is stored according to list B in a dry, dark place at a temperature from 5 °C to 25 °C. The shelf life is two years from the date of manufacture.

ENROGENT AQUA-TRV

enrofloxacin 10%
gentamicin 5%



Indications for use:

Enrogent aqua-TRV is used for therapeutic purposes in poultry with diseases of bacterial etiology, the pathogens of which are sensitive to the components of the drug.


Dosage: {  Poultry - 0,5-1 l/ton water

Storage conditions:

Enrogent aqua-TRV is stored under the requirement of the list B: in a dark place at temperatures from 5°C to 25°C. Shelf life is two years from the date of manufacture, subject to storage conditions.



ENROMOL AQUA-TRV

enrofloxacin 10%
colistin 2 000 000 IU/ml



Indications for use:

Enromol aqua-TRV is used as a therapeutic agent for treatment of respiratory and gastrointestinal infections of bacterial etiology, if the causative agents are sensitive to enrofloxacin and colistin.

Dosage: {  Poultry - 0,5-1 l/ton water
 Calves - 0,025-0,05 ml/kg of body weight

Storage conditions:

Enromol aqua-TRV is stored according to list B in a dry, dark place at a temperature of 5 °C to 25 °C. Shelf life is two years from the date of manufacture, subject to storage conditions.


ENROTRIM AQUA

enrofloxacin 10%
trimethoprim 5%



Indications for use:

Enrotrim aqua is used as a therapeutic agent in poultry (broiler chickens, replacement young laying hens, goslings, turkey poultry and ducklings) for treatment: colibacillosis, salmonellosis, necrotic enteritis, mycoplasmosis, etc.

Dosage: {  Poultry - 0,5 - 1 l/ton water

Storage conditions:

Enrotrim aqua is stored with precautions according to list B in a dry, dark place at a temperature from 5 °C to 25 °C. Shelf life is two years from the date of manufacture, subject to storage conditions.


FLOROL 10% AQUA-TRV

florfenicol 10%



Indications for use:

Florol 10% aqua-TRV is used as a therapeutic agent in poultry for treatment colibacillosis, salmonellosis, pasteurellosis, staphylococcosis and other infectious diseases of poultry caused by microorganisms susceptible to florfenicol.

Dosage: {  Poultry - 1-2 l/ton water

Storage conditions:

The veterinary drug is stored according to list B in a dry, dark place at a temperature from 5 °C to 25 °C. Shelf life - two years from the date of manufacture.


LESTIN AQUA

levofloxacin 10%
colistin 1 000 000 IU/ml



Indications for use:

The drug is used to treat poultry with bacterial infections of respiratory, digestive and genitourinary systems, in other bacterial infections, the pathogens of which are sensitive to components of the drug.

Dosage: {  Poultry - 0,4-0,5 l/ton water

Storage conditions:

The drug is stored in manufacturer's package according to the B-list in dry, protected from the sun place at temperature of 5°C up to 25°C. Shelf life – 24 months from the date of production under storage and transportation conditions.


LEXOL AQUA

levofloxacin 15%



Indications for use:

Lexol aqua is used for therapeutic purposes in poultry with bacterial infections, the pathogens of which are sensitive to levofloxacin.

Dosage: {  Poultry - 0,35-0,5 l/ton water

Storage conditions:

Lexol aqua is stored under the requirement of the list B: in a dark place at temperatures from 5°C to 25°C. Shelf life is two years from the date of manufacture, subject to storage and transportation conditions.



SULFAPRIM AQUA-TRV

sulfadimidin 20%
trimethoprim 4%



Indications for use:

Sulfaprim aqua-TRV is used for therapeutic purposes in cattle and poultry with diseases of bacterial etiology, the pathogens of which are sensitive to the components of the drug.

Dosage: {  Poultry - 0,25-1,0 l/ton water
 Calves - 1-2 ml/16 kg of body weight

Storage conditions:

Sulfaprim aqua-TRV is stored under the requirement of the list B: in a dark place at temperatures from 5°C to 25°C. Shelf life is three years from the date of manufacture, subject to storage conditions.


TETRAFLOX AQUA

tilmicosine 5%
levofloxacin 2,5%
trimethoprim 2,5%



Indications for use:

The preparation is administered for treatment of and farm poultry against diseases caused by microorganisms susceptible to the active substances of the preparation.

Dosage: {  Poultry - 1-2 l/ton water

Storage conditions:

The drug is stored in the manufacturer's packaging according to list B, in a dry, light-protected place at temperatures from plus 5 °C to plus 25 °C. Shelf life - two years from the date of manufacture.

TILMICOSIN 30% AQUA-TRV





tilmicosin 30%

Indications for use:

Tilmicosin 30% aqua-TRV is used as a therapeutic agent in poultry and calves with animal diseases, the causative agents of which are sensitive to tilmicosin.

Dosage:

-  Poultry - 250 ml/ton water
-  Cattle - 0,04 ml/kg of body weight

Storage conditions:

Tilmicosin 30% aqua-TRV is stored under the requirement of the list B: in a dark place at temperatures from 5°C to 25°C. Shelf life is three years from the date of manufacture, subject to storage and transportation conditions.

TRIENROKOL AQUA





**enrofloxacin 10%
trimethoprim 5%
colistin 1 000 000 IU/ml**

Indications for use:

Trienrokol aqua is used as a therapeutic agent for respiratory and gastrointestinal infections of bacterial etiology caused by microorganisms sensitive to components of the drug.

Dosage:

-  Poultry - 0,5 l/ton water
-  Calves - 0,25-0,5 ml/10 kg of body weight

Storage conditions:

Trienrokol aqua is stored under the requirement of the list B: in a dark place at temperatures from 5°C to 25°C. Shelf life is two years from the date of manufacture, subject to storage conditions.

TYLOSIN AQUA




tylosin 20%

Indications for use:

Tylosin aqua is used as a therapeutic agent: agricultural poultry (broilers, replacements, breeding poultry) - for respiratory mycoplasmosis, infectious sinusitis, colibacillosis, pasteurellosis and other diseases, pathogens of which are sensitive to tylosin.

Dosage:

-  Poultry - 2-3 l/ton water

Storage conditions:

Tylosin aqua is stored under the requirement of the list B: in a dark place at temperatures from 5°C to 25°C. Shelf life is two years from the date of manufacture, subject to storage conditions.



STOVEK
veterinary production

STOVEK LLC

222660, Republic of Belarus, Minsk region,
Stolbtsy, Zadvorjenskaya str., 2, office 67

8(017) 172 50 80,

e-mail: nemanbio@gmail.com

www.stovek.org




AMOXIDGEKT LA

amoxicillin 15%



Indications for use:

Amoxidgekt LA is used in cattle with infectious diseases of the gastrointestinal tract, respiratory system and genitourinary system; surgical diseases and diseases of the skin and soft tissues.

Dosage: {  Cattle - 1 ml/10 kg of body weight

Storage conditions:

Amoxidgekt LA is stored according to the requirements of the list B: in a dry, dark place at a temperature from 5 °C to 25 °C. The shelf life is two years from the date of manufacture, subject to the storage terms.



AZITHROMYCIN 10-TRV

azithromycin 10%



Indications for use:

Azithromycin 10% -TRV is used as a treatment for pigs, cattle, sheep in the treatment of bacterial infections caused by microorganisms sensitive to azithromycin.

Dosage: {  Cattle - 1 ml/20-40 kg of body weight
 Sheep - 1 ml/ 20kg of body weight

Storage conditions:

Azithromycin 10% -TRV is stored in a dry, dark place at temperatures from 5°C to 25°C. Shelf life - two years from the date of manufacture, subject to storage conditions.



CEFTIDGEKT LA-TRV

ceftiofur 10%



Indications for use:

The veterinary product is used for infectious diseases of bacterial etiology of the respiratory system, gastrointestinal tract and genitourinary system.

Dosage: {  Poultry - 0,05-0,2 ml/10 kg of body weight
 Calves - 1 ml/15 kg of body weight

Storage conditions:

Ceftidgekt LA-TRV is stored in a dry, dark place at temperatures from 5 °C to 25 °C. Shelf life is two years from the date of manufacture, provided that the storage conditions are observed.



ENRODGEKT 100

enrofloxacin 10%



Indications for use:

Enrodgekt 100 is used as a therapeutic agent for treatment colibacillosis, salmonellosis, pasteurellosis, pseudomonosis, clostridiosis, mycoplasmosis and other infectious diseases, pathogens are sensitive to enrofloxacin.

Dosage: {  Calves - 1 ml/40 kg of body weight
 lambs - 1 ml/40 kg of body weight

Storage conditions:

Enrodgekt 100 is stored in a dry, dark place at a temperature from 5 °C to 25 °C. Shelf life is two years from the date of manufacture, subject to storage conditions.

FLOREZOL



florfenicol 40%



Indications for use:

Florezol is used as a therapeutic agent in cattle and small stock for infectious diseases caused by pathogens sensitive to the action of the veterinary drug.

Dosage:

-  Calves - 0,75-1,5 ml/15 kg of body weight
-  Sheep - 0,75-1,5 ml/15 kg of body weight

Storage conditions:

Florezol is stored in a dry, dark place at temperatures from 5 °C to 25 °C. Shelf life is two years from the date of manufacture, subject to storage conditions.

GENTALUX


gentamicin 4%



Indications for use:

Gentalux is indicated for cattle with gastrointestinal, respiratory pathologies, sepsis, peritonitis, pyelonephritis, arthritis and other diseases of animals of bacterial etiology caused by microorganisms sensitive to antibiotics.

Dosage:

-  Cattle - 0,75-2 ml/10 kg of body weight

Storage conditions:

Gentalux is stored in a dry, dark place at temperatures from 5 °C to 25 °C. Shelf life - two years from the date of manufacture, subject to storage conditions.

LEXODGEKT



levofloxacin 15%



Indications for use:

The veterinary product is used for mixed and secondary bacterial infections, infectious diseases of bacterial etiology, the pathogens of which are sensitive to levofloxacin.

Dosage:

-  Cattle - 1 ml/30 kg of body weight
-  Sheep - 1 ml/ 30kg of body weight

Storage conditions:

Lexodgekt is stored according to list B in a dry, dark place at a temperature from 5 °C to 25 °C. The shelf life is two years from the date of manufacture, subject to the storage terms.

MOXIDGEKT LA-TRV


moxifloxacin 10%



Indications for use:

The drug is used for treatment of cattle with bacterial infections of the respiratory, digestive, genitourinary systems, infectious diseases of bacterial etiology the pathogens of which are sensitive to moxifloxacin.

Dosage:

-  Cattle - 1 ml/10 kg of body weight

Storage conditions:

The drug is stored in manufacturer's package according to the B-list in dry, protected from the sun place at temperature of 5°C up to 25°C. Shelf life – 24 months from the date of production.

OXSICET-TRV

oxytetracycline 20%
ketoprofen 3%



Indications for use:

Oxicet-TRV is used in cattle with combined infections, secondary bacterial infections in viral diseases.

Dosage:

 Cattle - 1 ml/10 kg of body weight

 Sheep - 1 ml/ 10kg of body weight

Storage conditions:

Oxicet-TRV is stored in a dry, dark place at temperatures from 5°C to 25°C. Shelf life is two years from the date of manufacture, subject to storage conditions.

OXYTETRACYCLINE 200 LA-TRV


oxytetracycline 20%



Indications for use:

Oxytetracycline 200 LA-TRV is used as a therapeutic agent for diseases of bacterial etiology, the causative agents of which are sensitive to oxytetracycline.

Dosage:

 Cattle - 1 ml/10 kg of body weight

 Sheep - 1 ml/ 10kg of body weight

Storage conditions:

Oxytetracycline 200 LA-TRV is stored according to list B in a dry, dark place at a temperature from 5 °C to 25 °C. The shelf life is two years from the date of manufacture, subject to the storage terms.

PENSTREP LA-TRV


benzylpenicillin procaine 10%
benzylpenicillin benzathine 10%
dihydrostreptomycin sulfate 20%




Indications for use:

The drug is used for treatment of cattle, sheep for respiratory diseases, gastrointestinal tract and urogenital system, musculoskeletal system and diseases caused by pathogens sensitive to the drug components.

Dosage:

 Cattle - 0,5-1 ml/10 kg of body weight

 Sheep - 0,5-1 ml/10 kg of body weight

Storage conditions:

The drug is stored in the manufacturer's package in List B, in a dry, light-protected place at temperatures from plus 2 °C to plus 15 °C. Shelf life - two years from the date of manufacture.

TULATHROMYCIN 10-TRV


tulathromycin 10%



Indications for use:

Tulathromycin 10-TRV is used in calves with bacterial respiratory infections (pasteurellosis, hemophillosis, mycoplasmosis), infectious keratoconjunctivitis and other diseases of bacterial etiology, the causative agents of which are sensitive to tulathromycin.

Dosage:

 Cattle - 1 ml/40 kg of body weight

Storage conditions:

Tulathromycin 10-TRV is stored according to the requirements of the list B: in a dry, dark place at a temperature from 5 °C to 25 °C. The shelf life is three years from the date of manufacture.

Injection antibiotics

TYLDGEKT



tylosin 20%



Indications for use:

Tyldgekt is used as a therapeutic agent for bronchopneumonia of cattle, bovine mastitis; arthritis, dysentery, as well as secondary bacterial diseases.

Dosage:

-  Cattle - 0,025-0,05 ml/kg of body weight
-  Sheep - 0,05-0,06 ml/kg of body weight

Storage conditions:

Tyldgekt is stored according to list B in a dry, dark place at a temperature from 5 °C to 25 °C. The shelf life is two years from the date of manufacture, subject to the storage terms.

Hormonal drugs

DECSA-TRV



dexamethasone 0,4%



Indications for use:

Decsa-TRV is administered to live-stock and domestic animals as anti-inflammatory, anti-allergic, decongestant and gluconeogenic agent in the treatment of post-traumatic oedema, arthritis, tenosynovitis.

Dosage:

-  Cattle - 4-9 ml/500 kg of body weight
-  Horses - 4-9 ml/ 500 kg of body weight

Storage conditions:

Decsa-TRV is stored in a dry, dark place at temperatures from 5 °C to 25 °C. Shelf life is two years from the date of manufacture, provided that the storage conditions are observed.

Hormonal drugs

CLOPROSTENOL-TRV


cloprostenol 0,025%



Indications for use:

Cloprostamol-TRV used in cows and calves for the regulation of reproductive function and treatment for gynecological and obstetric pathology, including synchronization of the sexual cycle in cows and calves; treatment of cows with functional disorders of the ovaries.

Dosage:

-  Cows - 2-4 ml/animal

Storage conditions:

Cloprostamol-TRV is stored under the requirement of the list B: in a dry, dark place at temperatures from 5°C to 25°C. Shelf life is two years from the date of manufacture, subject to storage conditions.

OXYTOCIN 10 ME-TRV



oxytocin 10 IU/ml



Indications for use:

Oxytocin 10 ME-TRV is used in agricultural and domestic animals with weak labor, uterine bleeding, retention of the placenta, endometritis, mastitis and reflexagalactia.

Dosage:

-  Cattle - 15-60 IU/animal
-  Sheep - 8-15 IU/animal

Storage conditions:

Oxytocin 10 ME-TRV is stored with caution (under the requirement of the list B): in a dry, dark place at temperatures from plus 5°C to plus 25°C. Shelf life is two years from the date of manufacture.

Anti-inflammatory drugs

KETOPROFEN 10-TRV





ketoprofen 10%

Indications for use:

Ketoprofen 10-TRV is used in cattle and horses as an anti-inflammatory, analgesic and antipyretic agent for acute and chronic diseases of the musculoskeletal system.

Dosage:

-  Cattle - 3 ml/100 kg of body weight
-  Horses - 1 ml/45 kg of body weight

Storage conditions:

Ketoprofen 10-TRV is stored in a dry, dark place at temperatures from plus 5°C to plus 25°C. Shelf life is two years from the date of manufacture, subject to storage conditions.

Metabolic stimulators

FERRUM 20/B12-TRV




iron 20%
vitamin B12 0,02%

Indications for use:

The veterinary drug is used for therapeutic and prophylactic purposes in calves, pigs, pregnant cows and heifers.

Dosage:

-  Calves - 2-4 ml/animal

Storage conditions:

The veterinary drug is stored in a dry, dark place at a temperature from 5 °C to 25 °C. Shelf life is two years from the date of manufacture, subject to storage conditions.

Antiparasitic drugs

IVERMECTIN 1%-TRV





ivermectin 1%

Indications for use:

The drug is used for therapeutic and prophylactic purposes in cattle with dictyocaulosis, haemonchosis, thelaziosis, strongyloidosis, esophagostomosis, bunostomosis, habertiosis, hypodermosis.

Dosage:

-  Cattle - 1 ml/50 kg of body weight
-  Sheep - 1 ml/50 kg of body weight

Storage conditions:

Ivermectin 1%-TRV is stored under requirements of list B in a dry, dark place at temperatures from 5°C to 25°C. Shelf life is three years from the date of manufacture, subject to storage conditions.

IVERMECTIN AQUA





ivermectin 4%

Indications for use:

The drug is administered to cattle, poultry, carnivores for prevention and treatment of nematodosis and arachnoentomosis.

Dosage:

-  Poultry - 100 ml/ton water
-  Calves - 0,25 ml/50 kg of body weight

Storage conditions:

The drug is stored in the manufacturer's packaging according to list B, in a dry, light-protected place at temperatures from plus 5 °C to plus 25 °C. Shelf life - two years from the date of manufacture.

Antiviral drugs

TRIVIRON FORTE-TRV



aRNase 3%



Indications for use:

Triviron forte-TRV is used for therapeutic and prophylactic purposes in animals and poultry with infections caused by RNA viruses.

Dosage:

-  Poultry - 0,00125-0,005 ml/kg of body weight
-  Calves - 0,05-0,1 ml/ kg of body weight

Storage conditions:

Triviron forte-TRV is stored according to the requirements of the list B: in a dry, dark place at a temperature from 4°C to 25 °C. The shelf life is two years from the date of manufacture, subject to the storage terms.

TRIVIDGEKT 5%-TRV


aRNase 5%



Indications for use:

Trividgekt 5%-TRV is used for therapeutic and prophylactic purposes in animals with infections caused by RNA viruses.

Dosage:

-  Calves -0,1-0,2 ml/kg of body weight

Storage conditions:

Trividgekt 5%-TRV is stored according to the requirements of the list B: in a dry, dark place at a temperature from plus 5 °C to 25 °C. The shelf life is two years from the date of manufacture, subject to the storage terms.

Feed additives

LYSINOFID-TRV


L-lysine 95%



Indications for use:

The feed additive is used as a source of L-lysine for enrichment and balancing the diets of farm animals, including poultry, preparation of complete feed, premixes and feed additives.

Dosage:

-  Poultry - 0.5 - 5% by weight of the finished feed

Storage conditions:

The additive is to be stored in manufacturer's package in dry, protected from light place at temperature of 0°C up to 25°C above zero. Shelf life – 24 months from the date of production.

TREOFID-TRV


L-threonine 95%



Indications for use:

The feed additive is used as a source of L-threonine for enriching and balancing diets of farm animals, including poultry, for preparation of complete feed mixes, premixes, feed additives.

Dosage:

-  Poultry - 1% by weight of the finished feed

Storage conditions:

Stored in the original manufacturer's packaging in dry, light-protected place at a temperature not exceeding 25 °C. Shelf life - 24 months from the date of production.

Probiotics

PROVAGEN

Animal Feed Supplement



**B.subtilis
B.licheniformis 10⁹ CFU/gram**

Indications for use:

Provagen is used in case of disruption of the functioning of the gastrointestinal tract against the background of disruption of natural microbiocenosis for the prevention of rumen acidosis and improving the digestibility of roughage in cattle, prevention of various stress effects, stimulation of immunity, increase in safety, decrease in feed conversion and increase in weight gain.

Dosage:

- Poultry - 0,1-1 kg/ton feed
- Calves - 0,1-1 kg/ton feed

Storage conditions:

Shelf life of the Supplement is 12 months from the date of manufacture, provided that it is stored in a dry, dark place at a temperature from 4 °C to 25 °C.

Probiotics

TRIOBAKT

**B.licheniformis
5x10⁹ CFU/gram**



Indications for use:

The additive is used for poultry as an adjuvant to normalise biocoenosis of the gut, prevention of stressful impact, as well as for stimulation of non-specific immunity and improvement of feed conversion.

Dosage:

- Poultry - 500 gram/ton feed
- Calves - 500 gram/ton feed

Storage conditions:

The additive is to be stored in manufacturer's package in dry, protected from light place at temperature of 0 °C up to 30 °C. Shelf life – 24 months from the date of production.

PROVAGEN CONCENTRATE

Animal Feed Supplement



**B.subtilis
B.licheniformis 10¹¹ CFU/gram**

Indications for use:

Supplement has a complex effect on the body, including: the competitive relationship of Bacillus subtilis and Bacillus licheniformis strains with pacified bacteria for the necessary nutrients or for adhesion; stimulation of immunity; antagonistic action of Bacillus subtilis and Bacillus licheniformis to both gram-positive and gram-negative pathogenic microorganisms.

Dosage:

- Poultry - 0,0002-0,001 gram/head
- Calves - 0,05-0,3 gram/animal

Storage conditions:

Shelf life of the supplement is 12 months from the date of manufacture, provided that it is stored in a dry, dark place at a temperature from 4 °C to 25 °C.

Biodestructor of manure

BIOSANVIT-M

Bacillus 10⁷ CFU/gram



Indications for use:

Complexed with bacteria of biodestructor "Biosanvit-M" act at temperature of + 4 °C to + 45 °C. The bacteria expansion starts in 12-16 hours. The bacteria become fully active in 2-4 days. The expected effect is observed starting from 6-10 days.

Dosage:

- Poultry - 50-70/gram ton of litter
- Calves - 150-200 gram/100 m² of area

Storage conditions:

Shelf life is 24 months from the date of manufacture, provided that it is stored at temperatures from 0 °C to 25 °C in a dry, dark place.

Antibiotic compatibility table

The interaction of antibiotics with simultaneous use can have both positive and negative results

++ gain the action
+ weak gain of the action
- weakening of the action
* increased toxicity

	Penicillins	Cephalosporins	Erythromycin	Tetracycline	Tylosin	Streptomycin	Neomycin	Kanamycin	Gentamicin	Lincomycin	Flumekvin	Sulfonamides	Tiamulin	Colistin	Spectinomycin	Doxycycline	Enrofloxacin
Penicillins		++	-	-	-	++	++	++	++	-	++	+	-	++	++	-	++
Cephalosporins	++		-	-	-	++	++	++	++	-	++	+	-	++	++	-	++
Erythromycin	-	-		++	++	-	++	-	-	+	*	++	++	++	++	++	*
Tetracycline	-	-	++		++	+	-	+	+	++	-	++	++	++	-	++	-
Tylosin	-	-	++	++		++	++	++	++	-	*	*	++	++	++	++	*
Streptomycin	++	++	-	+	++		*	*	*	+	++	+	++	*	++	-	++
Neomycin	++	++	++	-	++	*		*	*	+	++	+	++	*	++	-	++
Kanamycin	++	++	-	+	++	*	*		*	+	++	+	++	*	++	-	++
Gentamicin	++	++	-	+	++	*	*	*		-	++	+	++	*	++	-	++
Lincomycin	-	-	+	++	-	+	+	+	-		*	++	++	++	++	++	*
Flumekvin	++	++	*	-	*	++	++	++	++	*		+	++	++	++	-	++
Sulfonamides	+	+	++	++	*	+	+	+	-	++	+		++	++	-	-	+
Tiamulin	-	-	++	++	++	++	++	++	++	++	++	++		++	++	++	++
Colistin	++	++	++	++	++	*	*	*	*	++	++	++	++		*	++	++
Spectinomycin	++	++	++	-	++	++	++	++	++	++	++	-	++	*		-	++
Doxycycline	-	-	++	++	++	-	-	-	-	++	-	-	++	++	-		-
Enrofloxacin	++	++	*	-	*	++	++	++	++	*	++	+	++	++	++	-	



STOVEK LLC

222660, Republic of Belarus, Minsk region,
Stolbtsy, Zadvorienskaya str., 2, office 67

8(017) 172 50 80

e-mail: nemanbio@gmail.com

www.stovek.org

